Author index

Aalto, T. *see* Patiño, S.
Abbo, S., Lev-Yadun, S. & Galwey, N. Vernalization response of wild chickpea, 695
Agostini, D. *see* Vallorani, L.
Albaugh, T.J. *see* King, J.S.
Allegre, L. *see* Nogués, S.
Allen Jr, L.H. *see* Gesch, R.W.
Allen, H.L. *see* King, J.S.
Asensio, L. *see* Bordallo, J.J.
Ashford, A. Tubular vacuoles in arbuscular mycorrhizas, 545
Ashford, A.E. *see* Cairney, J.W.G.
Atzmon, N. *see* Shaul-Keinan, O.

Balasubramanian, S., Kim, S-J. & Podila, G.K. Differential expression of a malate synthase gene during the preinfection stage of symbiosis in the ectomycorrhizal fungus *Laccaria bicolor*, 517
Balduman, L. *see* Grulke, N.E.
Bangerth, F. *see* Clifton-Brown, J.C.
Bassman, J.H., Edwards, G.E. & Robberecht, R. Long-term exposure to enhanced UV-B radiation is not detrimental to growth and photosynthesis in Douglas-fir, 107
Baum, S. & Schwarze, F.W.M.R. Large-leaved lime (*Tilia platyphyllos*) has a low ability to compartmentalize decay fungi via reaction zone formation, 481
Belausov, E. *see* Shaul-Keinan, O.
Ben-Tal, Y. *see* Shaul-Keinan, O.
Bethke, P. Protocals for plant cell biologists, 13
Bliss, K.M., Jones, R.H., Mitchell, R.J. & Mou, P.P. Are competitive interactions influenced by spatial nutrient heterogeneity and root foraging behavior?, 409
Bonfante, P. *see* Novero, M.
Boote, K.J. *see* Gesch, R.W.
Bordallo, J.J., Lopez-Llorca, L.V., Jansson, H.-B., Salinas, J., Persmark, L. & Asensio, L. Colonization of plant roots by egg-parasitic and nematode-trapping fungi, 491
Bowes, G. *see* Gesch, R.W.
Bruhn, D. *see* Gavito, M.E.
Brundrett, M.C. Coevolution of roots and mycorrhizas of land plants, 275
Bryla, D. *see* Poulton, J.L.
Buford, M. *see* King, J.S.

Cairney, J.W.G. & Ashford, A.E. Biology of mycorrhizal associations of epacrids (Ericaceae), 305
Chaffey, N.J. Popular trees, specialist trees, 548
Chaignon, V., Di Malta, D. & Hinsinger, P. Fe-deficiency increases Cu acquisition by wheat cropped in a Cu-contaminated vineyard soil, 121
Charvet-Candela, V., Hitchin, S., Ernst, D., Sandermann Jr, H., Marmeisse, R. & Gay, G. Characterization of an *Aux/IAA* cDNA up-regulated in *Pinus pinaster* roots in response to colonization by the ectomycorrhizal fungus *Hebeloma cylindrosporum*, 769
Chet, I. *see* Shaul-Keinan, O.
Clifton-Brown, J.C., Lewandowski, I., Bangerth, F. & Jones, M.B. Comparative responses to water stress in stay-green, rapid- and slow senescing genotypes of the biomass crop, *Miscanthus*, 335
Cocucci, M. *see* Nocito, F.F.
Cojocariu, C. *see* Kreuzwieser, J.
Cotxarrera, L. *see* Nogués, S.

Davoli, P. *see* Weber, R.W.S.
Declerck, S. *see* Nielsen, J.S.
Dengate, S.B. *see* Renzaglia, K.S.
Di Malta, D. *see* Chaignon, V.
Díaz, S. *see* Vendramini, F.
Díaz, S., Mcintyre, S., Lavorel, S. & Pausas, J.G. Does hairiness matter in Harare? Resolving controversy in global comparisons of plant trait responses to ecosystem disturbance, 7
Dougherty, P. *see* King, J.S.
Duckett, J.G. *see* Renzaglia, K.S.
Duncan, G. *see* Glidewell, S.M.
Dyer, P.S. Hydrophobins in the lichen symbiosis, 1

Edwards, A.A. *see* Patiño, S.
Edwards, G.E. *see* Bassman, J.H.
Elad, Y. *see* Shaul-Keinan, O.
Engström, P. *see* Svensson, M.E.
Ernst, D. *see* Charvet-Candela, V.
Eshed, Y. *see* Shaul-Keinan, O.
Ezawa, T. *see* Uetake, Y.

Faccio, A. *see* Novero, M.
Fahey, T.J. *see* Lilleskov, E.
Fischer, B.U. *see* Suter, D.
Fotelli, M.N., Nahm, A., Heidenfelder, A., Papen, H., Rennenberg, H. & Geßler, A. Soluble nonprotein nitrogen compounds indicate changes in the nitrogen status of beech seedlings due to climate and thinning, 85
Frehner, M. *see* Suter, D.

Gadkar, V. *see* Shaul-Keinan, O.
Galwey, N. *see* Abbo, S.
Garbaye, J. *see* Jany, J.-L.
Garcia-Jacas, N. *see* Garnatje, T.
Gardes, M. An orchid–fungus marriage – physical promiscuity, conflict and cheating, 4
Garnatje, T., Vilatersana, R., Roché, C.T., Garcia-Jacas, N., Susanna, A. & Thill, D.C. Multiple introductions from the Iberian peninsula are responsible for the invasion of *Crupina vulgaris* in western North America, 419
Gartner, B.L. *see* Pruyn, M.L.
Gartner, B.L. *see* Spicer, R.
Gaut, B.D. Evolutionary dynamics of grass genomes, 15
Gavito, M.E., Bruhn, D. & Jakobsen, I. Phoshorous by arbuscular mycorrhizal hyphae does not increase when the host plant grows under atmospheric CO_2 enrichment, 751
Gay, G. *see* Charvet-Candela, V.
Genre, A. *see* Novero, M.

Gesch, R.W., Vu, J.C.V., Boote, K.J., Allen Jr, L.H. & Bowes, G. Sucrose-phosphate synthase activity in mature rice leaves following changes in growth CO_2 is unrelated to sucrose pool size, 77
Geßler, A. *see* Fotelli, M.N.
Ginzberg, I. *see* Shaul-Keinan, O.
Glidewell, S.M., Möller, M., Duncan, G., Mill, R.R., Masson, D. & Williamson, B. NMR imaging as a tool for noninvasive taxonomy: comparison of female cones of two Podocarpaceae, 197
Gobert, A. & Plassard, C. Differential NO_3^--dependent patterns of NO_3^- uptake in *Pinus pinaster, Rhizopogon roselus* and their ectomycorrhiza association, 509
Grace, J. *see* Patiño, S.
Griffin, K.L., Turnbull, M. & Murthy, R. Canopy position affects the temperature response of leaf respiration in *Populus deltoides*, 609
Groenhof, A.C. *see* Tilston, E.L.
Grulke, N.E., Preisler, H.K., Rose, C., Kirsch, J. & Balduman, L. O_3 uptake and drought stress effects on carbon acquisition on ponderosa pine in natural stands, 621
Grünzweig, J.M. *see* Shaul-Keinan, O.
Gurvich, D.E. *see* Vendramini, F.
Gyaneshwar, P., James, E.K., Reddy, P.M. & Ladha, J.K. *Herbaspirillum* colonization increases growth and nitrogen accumulation in aluminium-tolerant rice varieties, 131

Haisch, A. *see* Scherrer, S.
Harmon, M.E. *see* Pruyn, M.L.
Harris, P.J. *see* Trethewey, J.A.K.
Hartley-Whitaker, J. *see* Meharg, A.A.
Hartung, W. *see* Peuke, A.D.
Hautekèete, N.-C., Piquot, Y. & Van Dijk, H. Variations in ageing and meristematic activity in relation to flower-bud and fruit exclusion complex in the *Beta* species complex, 641
Heidenfelder, A. *see* Fotelli, M.N.
Hinsinger, P. *see* Chaignon, V.
Hitchin, S. *see* Charvet-Candela, V.
Hobbie, E.A. *see* Lilleskov, E.
Hodgson, J.G. *see* Vendramini, F.
Höfte, M. *see* Hulsen, K.
Högberg, M.N. & Högberg, P. Extramatrical ectomycorrhizal mycelium contributes one-third of the microbial biomass and produces, together with associated roots, half the dissolved organic carbon in a forest soil, 791
Högberg, P. *see* Högberg, M.N.
Hogetsu, t. *see* Zhou, Z.
Honegger, R. *see* Naef, A.
Honegger, R. *see* Scherrer, S.
Honegger, R. *see* Trembley, M.L.
Houle, G. The advantage of early flowering in the spring ephemeral annual plant *Floerkea proserpinacoides*, 689
Hulsen, K., Top, E.M. & Höfte, M. Biodegradation of linuron in a *Phaseolus* bioassay detected by chlorophyll fluorescence, 821

Ingles, P.J. *see* Møller, S.G.

Jakobsen, I. *see* Gavito, M.E.
Jakobsen, I. *see* Nielsen, J.S.
James, E.K. *see* Gyaneshwar, P.
Jansson, H.-B. *see* Bordallo, J.J.
Jany, J.-L., Garbaye, J. & Martin, F. *Cenococcum geophilum* populations show a high degree of genetic diversity in beech forests, 651
Joner, E.J. *see* Nielsen, J.S.
Jones, M.B. *see* Clifton-Brown, J.C.
Jones, R.H. *see* Bliss, K.M.
Jüssen, V. *see* Kreuzwieser, J.

Kaiser, R. *see* Naef, A.
Kapulnik, Y. *see* Shaul-Keinan, O.
Kim, S-J. *see* Balasubramanian, S.
King, J.S., Albaugh, T.J., Allen, H.L., Buford, M., Strain, B.R. & Dougherty, P. Below-ground carbon input to soil is controlled by nutrient availability and fine root dynamics in loblolly pine, 389
Kirsch, J. *see* Grulke, N.E.
Koide, R.T. *see* Poulton, J.L.
Kojima, T. *see* Uetake, Y.
Kouki, M. & Manetas, Y. Toughness is less important than chemical composition of *Arbutus* leaves in food selection by *Poecilimon* species, 399
Kranner, I. Gluthione status correlates with different degrees of desiccation tolerance in three lichens, 451
Kreuzwieser, J., Cojocariu, C., Jüssen, V. & Rennenberg, H. Elevated atmospheric CO_2 causes seasonal changes in carbonyl emissions from *Quercus ilex*, 327

Ladha, J.K. *see* Gyaneshwar, P.
Lamaze, L. *see* Pasche, F.
Lavorel, S. *see* Díaz, S.
Leake, J.R. *see* Mckendrick, S.L.
Lee Taylor, D. *see* Mckendrick, S.L.
Lev-Yadun, S. *see* Abbo, S.
Lewandowski, I. *see* Clifton-Brown, J.C.
Lilleskov, E., Hobbie, E.A. & Fahey, T.J. Ectomycorrizal fungal taxa differing in response to nitrogen deposition also differ in pure culture organic nitrogen use and natural abundance of nitrogen isotopes, 219
Lopez-Llorca, L.V. *see* Bordallo, J.J.
Lüscher, A. *see* Suter, D.

Manetas, Y. *see* Kouki, M.
Marmeisse, R. *see* Charvet-Candela, V.
Martin, F. *see* Jany, J.-L.
Masson, D. *see* Glidewell, S.M.
Mcintyre, S. *see* Díaz, S.
Mckendrick, S.L., Leake, J.R., Lee Taylor, D. & Read, D.J. Symbiotic germination and development of the myco-heterotrph *Neottie nidus-avis* in nature and its requirement for locally distributed *Sebacina* spp., 233
Meharg, A.A. & Hartley-Whitaker, J. Arsenic uptake and metabolism in resistant and nonresistant plant species, 29
Mill, R.R. *see* Glidewell, S.M.
Mitchell, R.J. *see* Bliss, K.M.
Möller, M. *see* Glidewell, S.M.
Møller, S.G., Ingles, P.J. & Whitelam, G.C. The cell biology of phytochrome signalling, 553

Mou, P.P. *see* Bliss, K.M.
Mulder, L. *see* Novero, M.
Murthy, R. *see* Griffin, K.L.

Naef, A., Roy, B.A., Kaiser, R. & Honegger, R. Insect-mediated reproduction of systemic infections by *Puccinia arrhenatheri* on *Berberis vulgaris*, 717
Nahm, A. *see* Fotelli, M.N.
Nielsen, J.S., Joner, E.J., Declerck, S., Olsson, S. & Jakobsen, I. Phospho-imaging as a tool for visualization and noninvasive measurement of P transport dynamics in arbuscular mycorrhizas, 809
Nocito, F.F., Sacchi, G.A. & Cocucci, M. Membrane depolarization induced K^+ efflux from subapical maize root segments, 45
Nogués, S., Cotxarrera, L., Allegre, L. & Trillas, M.I. Limitations to photosynthesis in tomato leaves induced by *Fusarium* wilt, 461
Nösberger, J. *see* Suter, D.
Novero, M., Faccio, A., Genre, A., Stougaard, J., Webb, K.J., Mulder, L., Parniske, M. & Bonfante, P. Dual requirement of the *LjSym4* gene for mycorrhizal development in epidermal and cortical cells of *Lotus japonicus* roots, 741

O'connor, P.J., Smith, S.E. & Smith, F.A. Arbuscular mycorrhizas influence plant diversity and community structure in a semiarid herbland, 209
Olsson, S. *see* Nielsen, J.S.

Papen, H. *see* Fotelli, M.N.
Parker, J.S , Rushing toward a phylogenetic future, 271
Parniske, M. *see* Novero, M.
Pasche, F., Pornon, A. & Lamaze, L. Do mature leaves provide a net source of nitrogen supporting shoot growth in *Rhododendron ferrugineum*?, 99
Pataki, D.E. Atmospheric CO_2 climate and evolution – lessons from the past, 10
Paterson, A.H. What has QTL mapping taught us about plant domestication?, 591
Patiño, S., Aalto, T., Edwards, A.A. & Grace, J. Is *Rafflesia* an endothermic plant?, 429
Paulitz, T.C. Biological webs of plant disease, 272
Pauls, K.P. *see* Schulze, D.
Pausas, J.G. *see* Díaz, S.
Persmark, L. *see* Bordallo, J.J.
Peuke, A.D., Schraml, C., Hartung, W. & Rennenberg, H. Identification of drought-sensitive beech ecotypes by physiological parameters, 373
Piccoli, G. *see* Vallorani, L.
Picon-Cochard, C. *see* Teyssonneyre, F.
Pierleoni, R. *see* Vallorani, L.
Piquot, Y. *see* Hautekéete, N.-C.
Pitt, D. *see* Tilston, E.L.
Plassard, C. *see* Gobert, A.
Podila, G.K. *see* Balasubramanian, S.
Podila, G.K. Signaling in mycorrhizal symbioses – elegant mutants lead the way, 541
Polidori, E. *see* Vallorani, L.
Pornon, A. *see* Pasche, F.
Poulton, J.L., Bryla, D., Koide, R.T. & Stephenson, A.G. Mycorrhizal infection and high soil phosphorus improve vegetative growth and the female and male functions in tomato, 255
Pregitzer, K.S. Fine roots of trees – a new perspective, 267
Preisler, H.K. *see* Grulke, N.E.
Pruyn, M.L., Gartner, B.L. & Harmon, M.E. Within-stem variation of respiration in *Psuedotsuga menziesii* (Douglas-fir) trees, 359

Rasmussen, H.N. & Whigham, D.F. Phenology of roots and mycorrhiza in orchid species differing in phototrophic strategy, 797
Read, D.J. *see* Mckendrick, S.L.
Reddy, P.M. *see* Gyaneshwar, P.
Rennenberg, H. *see* Fotelli, M.N.
Rennenberg, H. *see* Kreuzwieser, J.
Rennenberg, H. *see* Peuke, A.D.
Renzaglia, K.S., Dengate, S.B., Schmitt, S.J. & Duckett, J.G. Novel features of *Equisetum arvense* spermatozoids: insights into pteridophyte evolution, 159
Ringli, C. *see* Trembley, M.L.
Robberecht, R. *see* Bassman, J.H.
Roché, C.T. *see* Garnatje, T.
Rogers, H.J., An accessible root to modern plant science?, 14
Rose, C. *see* Grulke, N.E.
Roy, B.A. *see* Naef, A.
Rufty, T.W. *see* Tungate, K.D.

Sacchi, G.A. *see* Nocito, F.F.
Sacconi, C. *see* Vallorani, L.
Saito, M. *see* Uetake, Y.
Salinas, J. *see* Bordallo, J.J.
Sandermann Jr, H. *see* Charvet-Candela, V.
Schanz, F. *see* Walsby, A.E.
Scherrer, S., Haisch, A. & Honegger, R. Characterization and expression of *XPH1*, the hydrophobin gene of the lichen-forming ascomycete *Xanthoria parietina*, 175
Schmitt, S.J. *see* Renzaglia, K.S.
Schraml, C. *see* Peuke, A.D.
Schulze, D. & Pauls, K.P. Flow cytometric analysis of cellulose tracks development of embryogenic *Brassica* cells in microspore cultures, 249
Schwarze, F.W.M.R. *see* Baum, S.
Shaul-Keinan, O., Gadkar, V., Ginzberg, I., Grünzweig, J.M., Chet, I., Elad, Y., Wininger, S., Belausov, E., Eshed, Y., Atzmon, N., Ben-Tal, Y. & Kapulnik, Y. Hormone concentrations in tobacco roots change during arbuscular mycorrhizal colonization with *Glomus intraradices*, 501
Smilauer, P. *see* Smilauerová, M.
Smilauerová, M. & Smilauer, P. Morphological responses of plant roots to heterogeneity of soil resources, 703
Smith, F.A. *see* O'connor, P.J.
Smith, S.E. *see* O'connor, P.J.
Sochting, U. A new critical starting point for Antarctic lichen taxonomy, 550
Soussana, J.F. *see* Teyssonneyre, F.
Spicer, R. & Gartner, B.L. Compression wood has little impact on the water relations of Douglas-fir (*Psuedotsuga menziesii*) seedlings despite a large effect on shoot hydraulic properties, 633
Stephenson, A.G. *see* Poulton, J.L.
Stocchi, V. *see* Vallorani, L.
Stougaard, J. *see* Novero, M.

Strain, B.R. *see* King, J.S.
Susanna, A. *see* Garnatje, T.
Susko, D.J. *see* Tungate, K.D.
Suter, D., Frehner, M., Fischer, B.U., Nösberger, J. & Lüscher, A. Elevated CO_2 increases carbon allocation to the roots of *Lolium perenne* under free-air CO_2 enrichment but not in a controlled environment, 65
Svensson, M.E. & Engström, P. Closely related MADS-box genes in club moss (*Lycopodium*) show broad expression patterns and are structurally similar to, but phylogenetically distinct from, typical seed plant MADS-box genes, 439

Teyssonneyre, F., Picon-Cochard, C. & Soussana, J.F. How can we predict the effects of elevated CO_2 on the balance between perennial C_3 grass species competing for light?, 53
Thill, D.C. *see* Garnatje, T.
Thompson, K. *see* Vendramini, F.
Tilston, E.L., Pitt, D. & Groenhof, A.C. Composted recycled organic matter suppresses soil-borne diseases of field crops, 731
Top, E.M. *see* Hulsen, K.
Trembley, M.L., Ringli, C. & Honegger, R. Differential expression of hydrophobins *DGH1*, *DGH2* and *DGH3* and immunolocalization of DGH1 in strata of the lichenized basidocarp of *Dictyonema glabratum*, 185
Trethewey, J.A.K. & Harris, P.J. Location of (1 → 3)- and (1 → 3), (1 → 4)-β-D-glucans in vegetative cell walls of barley (*Hordeum vulgaris*) using immuno gold labelling, 347
Trillas, M.I. *see* Nogués, S.
Tungate, K.D., Susko, D.J. & Rufty, T.W. Reproduction and offspring competitiveness of *Senna obtusifolia* are influenced by nutrient availability, 661
Turnbull, M. *see* Griffin, K.L.

Uetake, Y., Kojima, T., Ezawa, T. & Saito, M. Extensive tubular vacuole system in an arbuscular mycorrhizal fungus, *Gigaspora margarita*, 761

Vallorani, L., Polidori, E., Sacconi, C., Agostini, D., Pierleoni, R., Piccoli, G., Zeppa, S. & Stocchi, V. Biochemical and molecular characterization of NADP-glutamate dehydrogenase from the ectomycorrhizal fungus *Tuber borchii*, 779
Van Dijk, H. *see* Hautekéete, N.-C.
Vendramini, F., Díaz, S., Gurvich, D.E., Wilson, P.J., Thompson, K. & Hodgson, J.G. Leaf traits as indicators of resource-use strategy in floras with succulent species, 147
Vilatersana, R. *see* Garnatje, T.
Vu, J.C.V. *see* Gesch, R.W.

Walsby, A.E. & Schanz, F. Light-dependent growth rate determines change in the population of *Planktothrix rubescens* over the annual cycle in Lake Zürich, Switzerland, 671
Webb, K.J. *see* Novero, M.
Weber, R.W.S. & Davoli, P. Autophagocytosis of carotenoid-rich lipid droplets into vacuoles during aeciospore ageing in *Puccinia distincta*, 471
Whigham, D.F. *see* Rasmussen, H.N.
Whitelam, G.C. *see* Møller, S.G.
Williamson, B. *see* Glidewell, S.M.
Wilson, P.J. *see* Vendramini, F.
Wininger, S. *see* Shaul-Keinan, O.

Zeppa, S. *see* Vallorani, L.
Zhou, Z. & Hogetsu, T. Subterranean community structure of ectomycorrhizal fungi under *Sullus grevillei* sporocarps in a *Lariix kaempferi* forest, 529

Keyword index

abscisic acid (ABA), 335, 373
acclimation, 77
acetaldehyde, 327
aeciospores, 471
Afrocarpus falcatus, 197
ageing, 641
allometry, 65
aluminium, 131
amino compounds, 85
antioxidants, 451
apoplast, 185
arbuscular mycorrhiza, 703, 761
arbuscular mycorrhizal (AM) fungi, 501, 541
arbuscular mycorrhizas, 209, 545
arsenic resistance, 29
arsenic, 29
Arthrobotrys oligospora, 491
assimilation, 621
Asteraceae, 419
astringency, 399
atmospheric CO_2, 10
atmospheric composition, 10
Australia, 305
autophagocytosis, 471
autoradiogram, 809
Aux/IAA, 769
auxin-regulated gene, 769
auxins, 501

basidiomycete, 517
beech (*Fagus sylvatica*), 373
beech, 85
benomyl, 209, 703
Beta species complex (wild beets), 641
bioassay, 821
biocontrol in soil systems, 731
biomass allocation, 661
biomass partitioning, 65
biomass, 335
bioremediation, 821
Biosphere 2, 609
Bird's nest orchid (Neottia nidus-avis), 4
Brassica napus, 249

C4 plants, 335
calcofluor white, 249
callose, 347
carbohydrates, 609
carbon allocation, 65
carbon dioxide, 77, 751
carbon, 131
carbonyl emission, 327
carotenoids, 471
carvacryl methyl ether, 717
catechin, 481
cell plates (1 → 3)-b-D-glucan (1 → 3),(1 → 4)-β-D-glucan, 347
cellulose, 249
Cenococcum geophilum, 219, 651
cereals, 731
chemotropism, 491
chickpea, 695
Chl*a* fluorescence, 821
chlorophyll fluorescence, 461
chlorophyll, 107
Cicer sp., 695
class 1 hydrophobin, 175
climate change, 53
climate, 85
CO_2, 429
coastal plain species, 409
colonization process, 741
colonization, 419, 751
community, 209
comparative ecology, 7, 147
comparative map, 15
competition for light, 53
competition, 529
compression wood, 633
condensed tannins, 399
conflicts of allocation, 689
copper, 121
Corallorhiza, 797
cortical block, 741
Cortinarius, 219
critical depth, 671
crop evolution, 695
Crupina vulgaris, 419
cyanobacteria, 671
cytokinin, 335
cytokinins, 501

deciduous forests, 689
degradation modes, 481
desiccation tolerance, 451
Dictyonema glabratum, 185
Dictyonema, 1
differential gene expression, 175
digital, 809
direct nitrogen transfers, 99
disease suppression, 731
disturbance, 7
diversity, 209
drought stress, 373, 621
drought, 335

earth system science, 10
ecosystem function, 10
ectomycorrhiza, 509, 517, 651
ectomycorrhizal (ECM) fungi, 219, 529, 791
ectomycorrhizal mycelium, 791
ectomycorrhizal symbiosis, 541
elevated CO_2, 65, 327
embryo development, 249
endocytosis, 471
endophytic growth, 491
Epacridaceae, 305
Equisetum, 159
Ericaceae, 305
ericoid mycorrhiza, 305
evenness, 209
evo-devo, 439
evolution, 4, 10, 275, 439
expression analysis, 779
extractable dissolved organic carbon (DOC), 791

ferns, 159
Festuca arundinacea, 53
fine root turnover, 389
fine roots, 267
fire, 7
fitness, 4, 255
flavanoids, 481
Floerkea proserpinacoides, 689
flow cytometry, 249
FLUENT™, 429
forest soil, 791
fragrances, 717
free-air CO_2 enrichment (FACE), 65
fruit set, 255
fumigation–extraction, 791
functional group, 209
functional types, 7
fungal specificity, 233
fungal viability, 741
fungi, 275

fungus, 1
Fusarium oxysporum, 461

Gaeumannomyces, 731
Galearis, 797
gallotannins, 399
gas exchange, 107
gene cloning, 779
gene family, 439
genetic diversity, 651
genome size, 15
gibberellins, 501
Gigaspora margarita, 761
global carbon cycle, 10
global change, 10
Glomus, 809
glutamate dehydrogenase, 779
glutathione, 451
glyoxylate pathway, 517
Goodyera, 797
grassland community, 703
grazing, 7
growing shoots, 99
growth rate, 671
growth, 107

hair roots, 305
Hebeloma cylindrosporum, 769
Herbaspirillum seropedicae, 131
herbivory, 399
heterotrophy, 797
Holcus lanatus, 53
Hordeum vulgare (barley), 347, 491
HPLC with post-column derivatization (HPLC-CRD), 481
hydraulic architecture, 633
hydraulic conductivity, 633
hydrophobicity, 1
hydrophobin, 1
hydrophobin, 185
hyperaccumulation, 29

immunogold labelling, 347
immunolocalization, 185
in situ germination, 233
in situ hybridisation, 1, 175, 185
indole, 717
inner bark, 359
internal transcribed spacer (ITS) polymorphism, 529
interspecific competition, 409
intron, 439
invasion founders, 419
iron, 121

jasmin lactone, 717

Laccaria bicolor, 219, 517
Lactarius theiogalus, 219
Lake Zürich, 671
land plants, 275
land use, 7
Larix kaempferi, 529
laser scanning confocal microscopy, 761
leaf development, 399
leaf thickness, 147
leaf toughness, 399
leaf water content, 147
lichen symbiosis, 175
lichen, 1
lichen-forming basidiomycete, 185
lichenized fruitbody, 185
lichens, 451
life-cycle gradient, 641
life-history, 689
linuron, 821
Liparis, 797
lipid droplets, 471
loblolly pine, 267, 389
Lolium perenne (ryegrass), 65
Lolium perenne, 53
low-temperature response loci, 695
Luzula campestris, 703
Lycopersicon esculentum (tomato), 255, 461), 491
Lycopsida, 439

MADS-box, 439
malate synthase gene, 517
malate synthase, 517
maternal effects, 661
meristematic activity, 641
metabolism, 29
metal acquisition, 121
methyl nicotinate, 717
microbial N, 791
microspore culture, 249
microsynteny, 15
mimicry, 429
minirhizotron, 389
Miscanthus, 335
mixture, 53
molecular markers, 419
monoculture, 53
monoxenic, 809
morphology, 1, 275
motile vacuoles, 545
MRI, 197
mutant, 769
mutants, 541
myco-heterotrohpy, 4, 233
mycorrhiza, 255, 751, 769
mycorrhizal mutualism, 4
mycorrhizal quantification, 741
mycorrhizal symbioses, 541
mycorrhizas, 275
mycotrophy, 797

^{15}N, 131
^{15}N labelling, 99
N remobilization, 85
N uptake, 85
net nitrogen source, 99
Nicotiana tabacum, 501
nitrate flux, 509
nitrate induction, 509
nitrogen fixation, 131
nitrogen isotopes, 219
nitrogen storage, 99
nitrogen uptake, 99
nitrogen, 609
NMR imaging, 197
NO_3^- selective microelectrode, 509
nutrient availability, 389
nutrient deficiency, 121
nutrient transfer checkpoints, 541
nutrients, 703
nutritional effects, 661

orchid, 233
Orchidaceae, 797
Oregon Green 488 carboxy-DFFDA, 761
organic nitrogen use, 219
organic phosphorus, 809
Oryza sativa, 131
osmoprotectants, 373
ozone (O_3) uptake, 621

papillae, 491
Phaseolus vulgaris (bean), 821
phenolics, 399
phenotypic variation, 29
Phoma, 731
phosphate, 809
phosphorus, 545
phosphorus, 751
photorespiration, 461
photosynthesis, 77, 107, 461, 621, 671
phylogeny, 15, 275, 439
phytohormones, 501
phytosiderophore, 121
Pinus pinaster, 509, 769
Pinus ponderosa (ponderosa pine), 621
Pinus resinosa, 517

Planktothrix rubescens, 671
plant cell walls, 347
plant functional types, 147
plant phylogeny, 159
plant traits, 7
Plantago lanceolata, 703
Plasmodiophora, 371
Poa angustifolia, 703
Poaceae, 347
podocarps, 197
pollen, 255
pollination, 429
pollution effects, 621
Populus deltoides (cottonwood), 609
potassium channels, 45
potassium efflux, 45
precision, 409
preinfection, 517
proanthocyanidins, 481
provenances, 373
Prumnopitys ferruginea, 197
Pseudotsuga menziesii (Douglas-fir), 359
Pseudotsuga menziesii var. *glauca*, 107
PSII-inhibitors, 821
pteridophytes, 159
Puccinia distincta, 471
^{14}C pulse labelling, 65
purification, 779

Q_{10}, 609
Quercus ilex, 327
Quercus pubescens, 327

Rafflesia, endothermy, 429
RAPD, 651
RAPDs, 419
reaction wood, 633
reaction zones, 481
relative growth rate (RGR), 689
reproductive schedule, 689
reproductive-parts excision, 641
resin canals, 197
resource allocation, 255
resource allocation, 641
resource-depletion hypothesis, 641
respiration, 429, 609
Rhizopogon roseolus, 509
Rhododendron ferrugineum, 99
rice, 77, 131
root colonization, 491
root dynamics, 389
root foraging behavior, 409
root morphology and topology, 703
roots, 275
root–shoot ratio, 65
rust fungi, 717

sapwood, 359
scale, 409
scaling, 609
scanning electron microscopy, 717
sclerophylly, 147
sclerotia, 651
seasonal dynamics, 267
seasonal phenology, 389
Sebacina spp., 4
Sebacina, 233
seed nitrogen, 661
seed size, 661
seedling development, 233
seedling performance, 661
senescence, 335, 641
Senna obtusifolia (sicklepod), 661
sensitivity, 409
signaling, 541
soil fertility, 255
soil microbial biomass, 791
soil nitrate availability, 85
soilborne pathogens, 731
spatial nutrient heterogeneity, 409
specific leaf area, 147
spermatia, 717
spermatozoids, 159
spring ephemeral annual plants, 689
stabilization of E_m, 45
stay-green, 335
stem aeration, 359
stem growth, 267
stem respiration, 359
stomatal conductance, 633
succulence, 147
sucrose-phosphate synthase, 77
Suillus grevillei, 529
summer cropping, 695
symbiosis, 1, 185
synteny, 15

taxonomy, 197
temperature, 609
terminal-RFLP (T-RFLP), 529
terrestrial ecosystems, 10
terrestrial orchids, 797
thinning, 85
Tipularia, 797
tobacco, 501
toxicity, 29
trade-off, 689
transcription factor, 439
transpiration, 373
transport, 545
Tricholoma inamoenum, 219
Tuber borchii, 779
tubular vacuoles, 545, 761

ultrastructure, 159
ultraviolet-B radiation (UV-B radiation), 107
underground dynamics, 529
uptake, 29
uptake, 751
UV-absorbing compounds, 107

vacuoles, 471
vascular wilt, 461
vegetative growth, 255
vernalization, 695
Verticillium chlamydosporium, 491
VOC, 327
volatile compounds, 429

water and nutrient availability, 267
water availability, 389
water potential, 373
weed reproduction, 661
wheat, 121

Xanthoria, 1
XPH1, 175

Zea mays, 45
zinc, 121